

Spectroscopy of Molecular Oxygen in the Atmosphere of Venus

F P Mills (Caltech, Pasadena, CA 91125; ph. 818-395-6447; e-mail: fpm@mercur1.gps.caltech.edu); D Crisp (JPL); S G Ryan (AAO);

Current Venusian atmospheric chemistry models [e.g., Sze & McElroy, *J. Planet. Space Sci.* (1975), Winick & Stewart, *JGR* (1980), Yung & DeMore, *Icarus* (1982), Krasnopolsky & Parshev, *Venus* (1983)] have identified chemical cycles that use H, Cl, and S compounds to catalyze the reformation of CO₂ from its photodissociation products (CO and O₂). These models predict that O₂ should be a minor, but detectable, constituent of the Venus atmosphere (column mixing ratio in the mesosphere of $\sim 1.5\text{--}3.5 \times 10^{-6}$ from the Yung & DeMore model). In addition, repeated observations since 1975 of intense O₂ airglow in the IR atmospheric bands near 1.269 μm have shown that ground-state O₂ is produced in the Venus nightside mesosphere at a rate comparable to that at which CO₂ is photodissociated on the dayside [Connes et al., *Ap. J.* (1979), Crisp et al., *JGR* (1996)]. Nevertheless, all attempts to measure O₂ abundances in the mesosphere have been unsuccessful. The current upper limit is 3×10^{-7} [Trauger and Lunine, *Icarus* (1983)].

Speculation has suggested that O₂ might have been depressed at the time of Trauger and Lunine's observations since SO₂ abundances declined by a factor of 8 from 1979-1988 [Esposito et al., *JGR* (1988), Na et al., *JGR* (1990)] and have since declined further [Na et al., DPS Conference (1995)]. If this were correct then O₂ might now be detectable.

We used the AAT's UHRF [Diego et al., *Mon. Not. R. Astr. Soc.* (1995)] to attempt to detect O₂ in the Venus mesosphere by observing the 763.6325 and 763.2165 nm lines at spectral resolving power ($\lambda/\Delta\lambda$) of 600,000 using the 1.5" square image slicer [Diego, *App. Opt.* (1993)].

Our preliminary analysis finds no detectable absorption from O₂ in the Venus mesosphere. Analysis of one morning's spectra gives a maximum line depth ($\Delta I/I$) of ≤ 0.005 (equivalent width I of $\sim 0.1 \text{ m}\text{\AA}$). Scaling from Trauger and Lunine's calculations gives a column mixing ratio of $\leq 3 \times 10^{-6}$, indicating the smaller SO₂ abundance may not have significantly affected the abundance of O₂ in the Venus mesosphere.

American Geophysical Union Abstract Form

Reference # 0000
Session 0.00

1. 1996 WPGM Meeting
2. No member or sponsor (1).
3. (a) Franklin P. Mills
Planetary Science
110-25 Caltech
Pasadena, CA 91125 USA
FPM@mercur1.gps.caltech.edu
(b) 818-395-6447
(c) 818-585-1917
4. SP
5. (i) S'lo
(1) 5405, 5464,
(")
6. P'
7. 0% published elsewhere
8. Charge \$50 to card, expires
9. C
10. Please schedule as poster.
Presenting author is also a
convenor of a parallel session.
11. Regular author

1 Date received: 8 FEB 96
Date formatted: March 15, 1996
Form version: 2.0